




Clean Version of Claims (9/20/02)

WE CLAIM:

1. An indole compound represented by the formula (I), or a pharmaceutically acceptable salt, solvate, or prodrug derivative thereof;



wherein ;

R₁ is selected from groups (a), (b), and (c) wherein;


(a) is C₇-C₂₀ alkyl, C₇-C₂₀ haloalkyl, C₇-C₂₀ alkenyl, C₇-C₂₀ alkynyl or carbocyclic radical, or

(b) is a member of (a) substituted with one or more independently selected non-interfering substituents; or



(c) is the group -(L₁)-R₁₁; where, -(L₁)- is a divalent linking group of 1 to 8 atoms and where R₁₁ is a group selected from (a) or (b);

R₂ is hydrogen, or a group containing 1 to 4 non-hydrogen atoms plus any required hydrogen atoms;


R_3 is $-(L_3)-Z$, where $-(L_3)-$ is a divalent linker group selected from a bond or a divalent group selected from:



and Z is selected from a group represented by the formulae,



or



wherein, X is oxygen and R_a is selected from hydrogen, C₁-C₈ alkyl, aryl, C₁-C₈ alkaryl, C₁-C₈ alkoxy, aralkyl and -CN;


R₄ is the group, -(L_C)-(acylamino acid group); wherein -(L_C)-, is an acylamino acid linker having an acylamino acid linker length of 1 to 8;

R₅ is selected from hydrogen or a non-interfering substituent;


R₆ and R₇ are selected from hydrogen or a non-interfering substituent.

2. The compound of claim 1 wherein R₂ is hydrogen, C₁-C₄ alkyl, C₂-C₄ alkenyl, -O-(C₁-C₃ alkyl), -S-(C₁-C₃ alkyl), C₃-C₄ cycloalkyl, -CF₃, halo, -NO₂, -CN, or -SO₃.

4. The compound of Claim 1 wherein the acylamino acid linker group, -(L_C)-, for R₄ is a divalent group selected from,




7. The compound of claim 1 wherein for R_3 , Z is the group represented by the formula;




and the linking group $-(L_3)-$ is a bond; and R_a is hydrogen, methyl, ethyl, propyl, isopropyl, phenyl or benzyl.

8. The compound of claim 1 wherein for R_3 , Z is the group represented by the formula;




and the linking group $-(L_3)-$ is a bond; and R_a is hydrogen.

9. The compound of claim 1 wherein for R_3 , Z is the group represented by the formula;




10. The compound of claim 1 wherein for R_3 , Z is the group represented by the formula;



11. The compound of Claim 1 wherein, for R₆ the non-interfering substituent is hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₇-C₁₂ aralkyl, C₇-C₁₂ alkaryl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkenyl, phenyl, tolulyl, xylenyl, biphenyl, C₁-C₈ alkoxy, C₂-C₈ alkenyloxy, C₂-C₈ alkynyloxy, C₂-C₁₂ alkoxyalkyl, C₂-C₁₂ alkoxyalkyloxy, C₂-C₁₂ alkylcarbonyl, C₂-C₁₂ alkylcarbonylamino, C₂-C₁₂ alkoxyamino, C₂-C₁₂ alkoxyaminocarbonyl, C₁-C₁₂ alkylamino, C₁-C₆ alkylthio, C₂-C₁₂ alkylthiocarbonyl, C₁-C₈ alkylsulfinyl, C₁-C₈ alkylsulfonyl, C₂-C₈ haloalkoxy, C₁-C₈ haloalkylsulfonyl, C₂-C₈ haloalkyl,


C₁-C₈ hydroxyalkyl, -C(O)O(C₁-C₈ alkyl), -(CH₂)_n-O-(C₁-C₈ alkyl), benzyloxy, phenoxy, phenylthio, -(CONHSO₂R), -CHO, amino, amidino, bromo, carbamyl, carboxyl, carbalkoxy, -(CH₂)_n-CO₂H, chloro, cyano, cyanoguanidinyl, fluoro, guanidino, hydrazide, hydrazino, hydrazido, hydroxy, hydroxyamino, iodo, nitro, phosphono, -SO₃H, thioacetal, thiocarbonyl, or carbonyl; where n is from 1 to 8.

12. The compound of Claim 1 wherein for R₁ the divalent linking group -(L₁)- is selected from a group represented by the formulae (Ia), (Ib), (Ic), (Id), (Ie), and (If):




13. The compound of claim 1 wherein the linking group -(L₁)- of R₁ is -(CH₂)-.

15. The compound of claim 1 wherein for R_1 the group R_{11} is a substituted or unsubstituted carbocyclic radical selected from the group consisting of cycloalkyl, cycloalkenyl, phenyl, spiro[5.5]undecanyl, naphthyl, norbornanyl, bicycloheptadienyl, tolulyl, xylenyl, indenyl, stilbenyl, terphenyl, diphenylethylenyl, phenyl-cyclohexenyl, acenaphthylenyl, and anthracenyl, biphenyl, bibenzyl and related bibenzyl homologues represented by the formula (a):



where n is a number from 1 to 8.

18. The compound of claim 1 wherein R_4 is the group, $-(L_C)-(acylamino\ acid\ group)$ and wherein the (acylamino acid group) is:



and R^{4a} is selected from the group consisting of H, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, heteroaryl and aryl; and wherein NR^{4b} is an amino acid residue with the nitrogen atom being part of the amino group of the amino acid.

21. A pharmaceutical formulation comprising a indole compound as claimed in claim 1 together with a pharmaceutically acceptable carrier or diluent therefor.

22. A method of inhibiting sPLA₂ mediated release of fatty acid which comprises contacting sPLA₂ with a therapeutically effective amount of indole compound as claimed in claim 1.

26. Use of a pharmaceutical composition comprising sPLA₂ inhibitor compounds according to Claim 1 and mixtures thereof for treatment of Inflammatory Diseases comprising administering a therapeutic amount of said compound to a patient in need thereof.